

ON THE ZERO SET OF THE KOBAYASHI–ROYDEN PSEUDOMETRIC OF THE SPECTRAL UNIT BALL

NIKOLAI NIKOLOV, PASCAL J. THOMAS

ABSTRACT. Given $A \in \Omega_n$, the n^2 -dimensional spectral unit ball, we show that if B is in the tangent cone C_A to the isospectral variety at A , then B is a "generalized" tangent vector at A to an entire curve in Ω_n if and only if B is in the tangent cone C_A to the isospectral variety at A . In the case of Ω_3 , the zero set of the Kobayashi–Royden pseudometric is completely described.

1. INTRODUCTION AND RESULTS

Let \mathcal{M}_n be the set of all $n \times n$ complex matrices. For $A \in \mathcal{M}_n$ denote by $sp(A)$ and $r(A) = \max_{\lambda \in sp(A)} |\lambda|$ the spectrum and the spectral radius of A , respectively. The spectral ball Ω_n is the set

$$\Omega_n := \{A \in \mathcal{M}_n : r(A) < 1\}.$$

The spectral Nevanlinna–Pick problem is the following one: given N points a_1, \dots, a_N in the unit disk $\mathbb{D} \subset \mathbb{C}$ and N matrices $A_1, \dots, A_N \in \Omega_n$ decide whether there is a mapping $\varphi \in \mathcal{O}(\mathbb{D}, \Omega_n)$ such that $\varphi(a_j) = A_j$, $1 \leq j \leq N$ (cf. [1, 2, 4, 7, 8] and the references there).

The study of the Nevanlinna–Pick problem in the case $N = 2$ reduces to the computation of the Lempert function, defined as follows for a domain $D \subset \mathbb{C}^m$:

$$l_D(z, w) := \inf\{|\alpha| : \exists \varphi \in \mathcal{O}(\mathbb{D}, D) : \varphi(0) = z, \varphi(\alpha) = w\}, \quad z, w \in D.$$

The infinitesimal version of the above is the Carathéodory–Fejér problem of order 1: given matrices $A_0, A_1 \in \mathcal{M}_n$, decide whether there is a mapping $\varphi \in \mathcal{O}(\mathbb{D}, \Omega_n)$ such that $A_0 = \varphi(0)$, $A_1 = \varphi'(0)$. This problem has been studied in [10].

2000 *Mathematics Subject Classification.* Primary: 32F45; Secondary: 32A07.

Key words and phrases. Spectral Nevanlinna–Pick problem, spectral Carathéodory–Fejér problem, spectral ball, symmetrized polydisc, Lempert function, Kobayashi–Royden pseudometric.

The initial version of this paper was written during the stay of the first named author at the Paul Sabatier University, Toulouse in May–June, 2007.

Its study reduces to the computation of the Kobayashi–Royden pseudometric, defined as follows for a domain $D \subset \mathbb{C}^m$:

$$k_D(z; X) := \inf\{|\alpha| : \exists \varphi \in \mathcal{O}(\mathbb{D}, D) : \varphi(0) = z, \alpha \varphi'(0) = X\}, \quad z \in D, X \in \mathbb{C}^m.$$

To each matrix A we associate its characteristic polynomial

$$P_A(t) := \det(tI - A) = t^n + \sum_{j=1}^n (-1)^j \sigma_j(A) t^{n-j},$$

where $I \in \mathcal{M}_n$ is the unit matrix,

$$\sigma_j(A) := \sigma_j(\lambda_1, \dots, \lambda_n) := \sum_{1 \leq k_1 < \dots < k_j \leq n} \lambda_{k_1} \dots \lambda_{k_j}$$

and $\lambda_1, \dots, \lambda_n$ are the eigenvalues of A .

Put $\sigma := (\sigma_1, \dots, \sigma_n) : \mathcal{M}_n \rightarrow \mathbb{C}^n$. The set

$$\mathbb{G}_n := \{\sigma(A) : A \in \Omega_n\}$$

is a taut (even hyperconvex) domain called the symmetrized n -disk (cf. [3, 8, 11] and references there). Explicit formulas for $l_{\mathbb{G}_2}$ and $k_{\mathbb{G}_2}$ can be found in [2] (see also [11]) and [10], respectively.

Recall now that a matrix $A \in \mathcal{M}_n$ is called non-derogatory if all the blocks in the Jordan form of A have distinct eigenvalues. Many equivalent properties to this definition may be found in [13, Proposition 3]. We point out one of them: A is non-derogatory if and only if $\text{rank}(\sigma_{*,A}) = n$, where $\sigma_{*,A}$ stands for the differential of σ at the point A .

Denote by \mathcal{C}_n the open and dense set of all non-derogatory matrices in Ω_n .

If $A_1, \dots, A_N \in \mathcal{C}_n$, then any mapping $\varphi \in \mathcal{O}(\mathbb{D}, \mathbb{G}_n)$ with $\varphi(\alpha_j) = \sigma(A_j)$ can be lifted to a mapping $\tilde{\varphi} \in \mathcal{O}(\mathbb{D}, \Omega_n)$ with $\tilde{\varphi}(\alpha_j) = A_j$, $1 \leq j \leq N$ [1]. This means that in a generic case the spectral Nevanlinna–Pick problem for Ω_n (with dimension n^2) can be reduced to the standard Nevanlinna–Pick problem for \mathbb{G}_n (with dimension n).

By the contractibility of the Lempert function, we have

$$l_{\Omega_n}(A, B) \geq l_{\mathbb{G}_n}(\sigma(A), \sigma(B)), \quad A, B \in \Omega_n,$$

and the lifting above implies that equality holds when both $A, B \in \mathcal{C}_n$.

By this and the fact that \mathbb{G}_n is a bounded domain, $l_{\Omega_n}(A, B) > 0$ if $sp(A) \neq sp(B)$. On the other hand, if $sp(A) = sp(B)$, then there is an entire mapping $\varphi : \mathbb{C} \rightarrow \Omega_n$ with $\varphi(0) = A$ and $\varphi(1) = B$ [9]; note that $sp(\varphi(\zeta)) = sp(A)$ for all $\zeta \in \mathbb{C}$, since by Liouville's Theorem, whenever $\varphi(\mathbb{C}) \subset \Omega$, then $\sigma \circ \varphi$ is constant. This situation is similar

to that of Brody's theorem for compact manifolds [5]: failure of hyperbolicity (that is, vanishing of the pseudodistance) can be explained by the presence of a (nonconstant) entire curve in the manifold.

Restricting again to non-derogatory matrices, a similar lifting [10] implies in the Carathéodory-Fejér case

$$k_{\Omega_n}(A; B) = k_{\mathbb{G}_n}(A, \sigma_{*,A}(B)), \quad A \in \mathcal{C}_n, B \in \mathcal{M}_n;$$

in particular, $k_{\Omega_n}(A; B) = 0$ if and only if $\sigma_{*,A}(B) = 0$. On the other hand, if $\sigma_{*,A}(B) = 0$, then there is an entire mapping $\varphi : \mathbb{C} \rightarrow \Omega_n$ with $\varphi(0) = A$ and $\varphi'(0) = B$ (indeed, $B = [Y, A] := YA - AY$ for some $Y \in \mathcal{M}_n$ [13, Proof of Proposition 3] and then the mapping $\zeta \rightarrow e^{\zeta Y} A e^{-\zeta Y}$ does the job).

The aim of this paper is to study the zeros of $B \rightarrow k_{\Omega_n}(A; B)$ in the remaining case, where A is a derogatory matrix, and to relate it to the existence of entire curves tangent to B at the point A (which is an obvious sufficient condition for $k_{\Omega_n}(A; B) = 0$).

For $A \in \Omega_n$ denote by C_A the tangent cone (cf. [6, p. 79] for this notion) to the isospectral variety

$$L_A := \{C \in \Omega_n : sp(C) = sp(A)\},$$

that is

$$C_A := \{B \in \mathcal{M}_n : \exists 0 < c_j \rightarrow 0, C_j \in L_A \text{ with } c_j(C_j - A) \rightarrow B\}.$$

Observe that L_A is smooth at A if $A \in \mathcal{C}_n$; then $C_A = \ker \sigma_{*,A}$. When $A \notin \mathcal{C}_n$, the rank of $\sigma_{*,A}$ is not maximal, so we have $\dim \ker \sigma_{*,A} > n^2 - n$; by [6, Corollary, p. 83], C_A is an analytic set with $\dim C_A = \dim L_A = n^2 - n$, so we have $C_A \subsetneq \ker \sigma_{*,A}$.

The following proposition characterizes the tangent cone C_A as the set of "generalized" tangent vectors at A to an entire curve in Ω_n through A (therefore contained in L_A).

Proposition 1. *Let $A \in \Omega_n$ and $B \in \mathcal{M}_n$. Then there are $m \in \mathbb{N}^1$, $m \leq n!$, and $\varphi \in \mathcal{O}(\mathbb{C}, \Omega_n)$ with $\varphi(0) = A$, $\varphi'(0) = \dots = \varphi^{(m-1)}(0) = 0$, $\varphi^{(m)}(0) = B$ if and only if $B \in C_A$.*

Proposition 1 implies that C_A is contained in the zero set of the singular Kobayashi pseudometric (cf. [14])

$$\hat{k}_{\Omega_n}(A; B) = \inf\{|\alpha| : \exists m \in \mathbb{N}, \varphi \in \mathcal{O}(\mathbb{D}, \Omega_n) : \text{ord}_0(\varphi - z) \geq m, \alpha \varphi^{(m)}(0) = m!X\}.$$

A consequence of the proof of Proposition 1 is the following.

¹ $\mathbb{N} := \{m \in \mathbb{Z} : m > 0\}$

Corollary 2. *Let $A \in \Omega_n$ and $B \in C_A$. Then the following conditions are equivalent:*

- (a) *there is $\varphi \in \mathcal{O}(\mathbb{C}, \Omega_n)$ with $\varphi(0) = A$ and $\varphi'(0) = B$.*
- (b) *there are $(r_j) \rightarrow \infty$ and $\varphi_j \in \mathcal{O}(r_j \mathbb{D}, \Omega_n)$, $j \in \mathbb{N}$, uniformly bounded near 0, such that $\varphi_j(0) = A$ and $\varphi_j'(0) = B$.*
- (c) *there are $r > 0$ and $\varphi \in \mathcal{O}(r \mathbb{D}, L_A)$ with $\varphi(0) = A$ and $\varphi'(0) = B$.*

Before stating the next proposition, we shall define an algebraic cone $C'_A \subset \mathcal{M}_n$, $A \in \Omega_n$.

For a function g holomorphic near A , and X in a neighborhood of A , let $g(X) - g(A) = g_A^*(X - A) + \dots$, where g_A^* stands for the homogeneous polynomial of lowest nonzero degree in the expansion of g near A (the omitted terms are thus of higher order).

Set

$$C_A^* := \{B \in \mathcal{M}_n : (\sigma_1)_A^*(B) = 0, \dots, (\sigma_n)_A^*(B) = 0\},$$

$$C'_A := \bigcap_{\lambda \in sp(A)} (\Phi_\lambda)_{*,A}^{-1}(C_{\Phi_\lambda(A)}^*),$$

where

$$(1) \quad \Phi_\lambda(A) := (A - \lambda I)(I - \bar{\lambda}A)^{-1}.$$

Note that

$$C_A \subset C_A^* \subset \ker \sigma_{*,A}.$$

For the first inclusion, cf. [6, p. 86, lines 4–6]), and for the second one, use that

$$\ker(\sigma_{*,A}) = \{(\sigma_j)_A^* = 0 \text{ for all } j \text{ such that } \deg(\sigma_j)_A^* = 1\}.$$

Since C_A and $\ker \sigma_{*,A}$ are invariant under automorphisms of Ω_n , it follows that

$$C_A \subset C'_A \subset \ker \sigma_{*,A}.$$

Moreover, if $\dim C_A = \dim C_A^*$, that is, $\dim C_A^* = n^2 - n$, then $C_A = C_A^* = C'_A$ (cf. [6, p. 112, Corollary 2]).

Proposition 3. *Let $A \in \Omega_n \setminus \mathcal{C}_n$.*

- (i) *If $\hat{k}_{\Omega_n}(A; B) = 0$, then $B \in C'_A$;*
- (ii) *$C'_A \neq \ker \sigma_{*,A}$.*

Remark. The cone C_A^* may coincide with $\ker \sigma_{*,A}$ for some $A \in \Omega_n \setminus \mathcal{C}_n$, $n \geq 3$. For example, if $A := \text{diag}(t, \dots, t, 0)$, $t \in \mathbb{D}_*$, then

$$C_A^* = \ker \sigma_{*,A} = \{B \in \mathcal{M}_n : \text{tr} B = b_{nn} = 0\}.$$

The main consequence of Proposition 3 is the fact that for $A \in \Omega_n \setminus \mathcal{C}_n$ and $B \in \ker \sigma_{*,A} \setminus C'_A$, a lifting for the corresponding Carathéodory-Fejér problem is not possible and $k_{\Omega_n}(\cdot; B)$ is not a continuous function

at A . This generalizes previous discontinuity results (see [13] and references therein).

Note also that the cone C'_A may coincide with C_A in some cases, for example, for any $A \in \Omega_2$ (then $C_A^* = C_A$, too) and any $A \in \Omega_3$ (for the last, see Proposition 6 and the discussion before it). We do not know whether this holds in general. On the other hand, it is not hard to find cases where $C_A \subsetneq C_A^*$.

Proposition 4. *For any $n \geq 3$ there is $A \in \Omega_n$ such that $C_A \subsetneq C_A^*$.*

Now, we state a conjecture about the zero set of k_{Ω_n} .

Conjecture 5. *$k_{\Omega_n}(A; B) = 0$ if and only if there is $\varphi \in \mathcal{O}(\mathbb{C}, \Omega_n)$ with $\varphi(0) = A$ and $\varphi'(0) = B$. In particular, if $k_{\Omega_n}(A; B) = 0$, then $B \in C_A$.*

Conversely however, there are matrices $B \in C_A$ such that $k_{\Omega_n}(A; B) \neq 0$ (see Proposition 6(ii) and Corollary 7).

There are some cases where our conjecture can be checked.

For example, since Ω_n is a balanced domain, one has that $l_{\Omega_n}(0, \cdot)$ and $k_{\Omega_n}(0; \cdot)$ coincide with the Minkowski function, that is, with the spectral radius. Thus the zeros of $k_{\Omega_n}(0; \cdot)$ are exactly the zero-spectrum matrices, and the set of those matrices is a union of complex lines through 0.

Also, if A is a scalar matrix, that is, $A = \lambda I$, $\lambda \in \mathbb{C}$, then $B \in C_A$, if and only if there is $\varphi \in \mathcal{O}(\mathbb{C}, \Omega_n)$ with $\varphi(0) = A$ and $\varphi'(0) = B$. To see this, use an automorphism of Ω_n of the form (1) to reduce to the case $A = 0$.

Since the derogatory matrices in Ω_2 are exactly the scalar matrices, we may choose $m = 1$ in Proposition 1 if $n = 2$, and C_A coincides both with the zeros of $k_{\Omega_2}(A; \cdot)$ and with the matrices $B = \varphi'(0)$ for some entire curve φ in Ω_2 (on the other hand, $\ker \sigma_{*,A} = \{B \in \mathcal{M}_2 : \text{tr} B = 0\}$).

Now we shall study the zero set of $k_{\Omega_3}(A; \cdot)$, when A is a non-scalar derogatory matrix. Using first an automorphism of the form (1) and then an automorphism of the form $C \rightarrow D^{-1}CD$, reduces the problem to the following two cases:

$$A = A_t := \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & t \end{pmatrix}, \quad t \in \mathbb{D}_*, \quad A = \tilde{A} := \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix}.$$

It is easy to see that

$$C_{A_t} \subset C_{A_t}^* = C'_{A_t} = \{B \in \mathcal{M}_3 : b_{33} = b_{11} + b_{22} = b_{11}^2 + b_{12}b_{21} = 0\},$$

$$C_{\tilde{A}} \subset C'_{\tilde{A}} = \{B \in \mathcal{M}_3 : b_{11} + b_{22} + b_{33} = b_{32} = b_{12}b_{31} = 0\}$$

(to prove, for example, the second inclusion, use that if $B_\varepsilon = A + \varepsilon B + o(\varepsilon)$, then $\text{tr} B_\varepsilon = \varepsilon \text{tr} B + o(\varepsilon)$, $\sigma_2(B_\varepsilon) = -\varepsilon b_{32} + o(\varepsilon)$ and $\det B_\varepsilon = \varepsilon^2(b_{12}b_{31} - b_{11}b_{32}) + o(\varepsilon^2)$). The next proposition implies, in particular, that $C_{A_\lambda} = C'_{A_\lambda}$ and $C_{\tilde{A}} = C'_{\tilde{A}}$ (use the fact that the tangent cones are closed or the dimensional reasoning mentioned above).

Proposition 6. (i) For any $B \in C'_{A_t}$ ($t \neq 0$) there is $\varphi \in \mathcal{O}(\mathbb{C}, \Omega_3)$ with $\varphi(0) = A_t$ and $\varphi'(0) = B$.

(ii) Let $B \in C'_{\tilde{A}}$. Then there is $\varphi \in \mathcal{O}(\mathbb{C}, \Omega_n)$ with $\varphi(0) = \tilde{A}$ and $\varphi'(0) = B$ if and only if $b_{11} = 0$ or $b_{12} \neq b_{31}$. Otherwise, $k_{\Omega_3}(\tilde{A}; B) > 0$.

Corollary 7. For any $n \geq 3$ there are $A \in \Omega_n$ and $B \in C_A$ such that $k_{\Omega_n}(A; B) > 0$.

Since $k_{\Omega_3}(A; B) > 0$ if $B \notin C'_A$, Proposition 6 and the discussion before give a complete description of the zero set of k_{Ω_3} .

Note that the situation is much easier for the Carathéodory-Reiffen pseudometric

$$\gamma_{\Omega_n}(A; B) = \sup\{|f'(A)B| : f \in \mathcal{O}(D, \mathbb{D})\}.$$

Here $\gamma_{\Omega_n}(A; B) = 0$ if and only if $\sigma_{*,A}(B) = 0$. Indeed, if $\sigma_{*,A}(B) \neq 0$, then

$$\gamma_{\Omega_n}(A; B) \geq \gamma_{\mathbb{G}_n}(A; \sigma_{*,A}(B)) > 0.$$

On the other hand, if $A \in \mathcal{C}_n$ and $\sigma_{*,A}(B) = 0$, then

$$0 = k_{\Omega_n}(A; B) \geq \gamma_{\Omega_n}(A; B) \geq 0.$$

It remains to use the density of \mathcal{C}_n in Ω_n and the continuity of the Carathéodory-Reiffen pseudometric.

The rest of the paper is organized as follows. The proofs of Propositions 3 and 4 are given in Section 2, the proofs of Proposition 6 and Corollary 7 – in Section 3, and the proofs of Proposition 1 and Corollary 2 – in Section 4.

Acknowledgment. The authors wish to thank the referee for a quick, thorough and perceptive report, thanks to which several mistakes in the first version have been corrected.

2. PROOFS OF PROPOSITIONS 3 AND 4

Proof of Proposition 4. Set $A := \text{diag}(0, \dots, 0, t, t)$, $t \in \mathbb{D}_*$. It is easy to see that $(\sigma_1)_A^*(B) = \sum_{j=1}^n b_{jj}$, $(\sigma_2)_A^*(B) = 2t \sum_{j=1}^{n-2} b_{jj} + t(b_{n-1, n-1} + b_{nn})$, $(\sigma_3)_A^*(B) = t^2 \sum_{j=1}^{n-2} b_{jj}$. Therefore, $(\sigma_3)_A^* = t(\sigma_2)_A^* - t^2(\sigma_1)_A^*$ and $\dim C_A^* > n^2 - n = \dim C_A$.

Proof of Proposition 3. (i) Let $\hat{\gamma}_{\Omega_n}(A; B)$ be the singular Carathéodory metric (cf. [12])

$$\hat{\gamma}_{\Omega_n}(A; B) := \sup \left\{ \left| \frac{f^{(k)}(A)B}{k!} \right|^{1/k} : k \in \mathbb{N}, f \in \mathcal{O}(\Omega_n, \mathbb{D}), \text{ord}_A f \geq k \right\},$$

where $\left| \frac{f^{(k)}(A)B}{k!} \right| = \sum_{|\alpha|=k} D^\alpha f(A) B^\alpha$. Since

$$\hat{k}_{\Omega_n}(A; B) \geq \hat{\gamma}_{\Omega_n}(A; B),$$

it is enough to show that $\hat{\gamma}_{\Omega_n}(A; B) > 0$ if $B \notin C'_A$. Then $B \in C_{\Phi_\lambda(A)}^*$ for some $\lambda \in sp(A)$. Replacing A and B by $\Phi_\lambda(A)$ and $(\Phi_\lambda)_{*,A}(B)$, respectively, we may assume that $B \notin C_A^*$. Then there is σ_j such that $(\sigma_j)_A^*(B) \neq 0$. Denoting by k the degree of $(\sigma_j)_A^*$, it follows that

$$\hat{\gamma}_{\Omega_n}(A; B) \geq \left| \frac{(\sigma_j)_A^*(B)}{\binom{n}{j}} \right|^{\frac{1}{k}} > 0.$$

(ii) Since $A \in \Omega_n \setminus \mathcal{C}_n$, at least two of the eigenvalues of A are equal, say to λ . Applying the automorphism Φ_λ of Ω_n , we may assume that $\lambda = 0$. Since the map $A \rightarrow P^{-1}AP$ is a linear automorphism of Ω_n for any $P \in \mathcal{M}_n^{-1}$, we may also assume that A is in Jordan form. In particular,

$$A = \begin{pmatrix} A_0 & 0 \\ 0 & A_1 \end{pmatrix},$$

where $A_0 \in \mathcal{M}_m$, $2 \leq m \leq n$, $sp(A_0) = \{0\}$, $A_1 \in \mathcal{M}_{n-m}$, $0 \notin sp(A_1)$. Furthermore, there is a set $J \subsetneq \{2, \dots, m\}$, possibly empty, such that $a_{j-1,j} = 1$ for $j \in J$, and all other coefficients $a_{ij} = 0$ for $1 \leq i, j \leq m$. Denote $0 \leq r := \#J = \text{rank} A_0 \leq m - 2$.

We set $B := \begin{pmatrix} B_0 & 0 \\ 0 & 0 \end{pmatrix} \in \mathcal{M}_n$, where $B_0 = (b_{ij})_{1 \leq i, j \leq m}$ is such that $b_{j-1,j} = -1$ for $j \in \{2, \dots, m\} \setminus J$, $b_{m1} = 1$, and $b_{ij} = 0$ otherwise. To complete the proof, it is enough to show the following.

Lemma 8. $(\sigma_m)_A^*(B) = 1$, but $\sigma_{*,A}(B) = 0$.

Proof. We begin by computing $\sigma_j(A_0 + hB_0)$, $1 \leq j \leq m$, $h \in \mathbb{C}$. Expanding with respect to the first column, we see that

$$\det(tI - (A_0 + hB_0)) = t^m + (-1)^{m-1} h^{m-r}.$$

Comparing the corresponding coefficients of both sides, it follows that

$$(2) \quad \sigma_j(A_0 + hB_0) = \begin{cases} 0, & 1 \leq j \leq m-1 \\ h^{m-r}, & j = m \end{cases}.$$

Next, we need a general formula for the functions σ_j . Given a matrix $M = (m_{ij})_{1 \leq i, j \leq n}$ and a set $E \subset \{1, \dots, n\}$, we write $\delta_E(M)$ for the determinant of the matrix $(m_{ij})_{i, j \in E} \in \mathcal{M}_{\#E}$. By convention, $\delta_\emptyset(M) = \sigma_0(M) := 1$. Then

$$(3) \quad \sigma_j(M) = \sum_{E \subset \{1, \dots, n\}, \#E=j} \delta_E(M).$$

Because of the block structure of our matrices,

$$\delta_E(A + hB) = \delta_{E \cap \{1, \dots, m\}}(A_0 + hB_0) \delta_{E \cap \{m+1, \dots, n\}}(A_1).$$

Therefore

$$\begin{aligned} \sigma_j(A + hB) &= \sum_{\max(0, j-n+m) \leq k \leq \min(m, j)} \left(\sum_{E' \subset \{1, \dots, m\}, \#E'=k} \delta_{E'}(A_0 + hB_0) \right) \times \\ &\quad \times \left(\sum_{E'' \subset \{m+1, \dots, n\}, \#E''=j-k} \delta_{E''}(A_1) \right) = \\ &\quad \sum_{\max(0, j-n+m) \leq k \leq \min(m, j)} \sigma_k(A_0 + hB_0) \sigma_{j-k}(A_1). \end{aligned}$$

It follows by (2) that $\sigma_j(A + hB) = S_1 + S_2$, where

$$S_1 = \begin{cases} \sigma_j(A_1), & j \leq n - m \\ 0, & \text{otherwise} \end{cases}, \quad S_2 = \begin{cases} h^{m-r} \sigma_{j-m}(A_1), & j \geq m \\ 0, & \text{otherwise} \end{cases}.$$

In particular,

$$\sigma_j(A) = \begin{cases} \sigma_j(A_1), & j \leq n - m \\ 0, & \text{otherwise} \end{cases}.$$

Then

$$\sigma_j(A + hB) - \sigma_j(A) = \begin{cases} h^{m-r} \sigma_{j-m}(A_1), & j \geq m \\ 0, & \text{otherwise} \end{cases}$$

Since $m - r \geq 2$ we get that $\sigma_{*,A}(B) = 0$, but $(\sigma_m)_A^*(B) = 1$. \square

3. PROOFS OF PROPOSITION 6 AND COROLLARY 7

Proof of Proposition 6. (i) Let first $B \in C'_{A_t}$. We shall write B in the form $B = X + [Y, A_t]$, where X is such that $\psi(\zeta) = A_t + \zeta X \in L_{A_t}$ for any $\zeta \in \mathbb{C}$. Then $\varphi(\zeta) = e^{\zeta Y} \psi(\zeta) e^{-\zeta Y}$ has the required properties.

It is easy to compute that $\psi(\mathbb{C}) \subset L_{A_t}$ if and only if $sp(X) = 0$ and $x_{11} + x_{22} = x_{11}^2 + x_{12}x_{21} = 0$. On the other hand,

$$[Y, A_t] = t \begin{pmatrix} 0 & 0 & y_{13} \\ 0 & 0 & y_{23} \\ -y_{31} & -y_{32} & 0 \end{pmatrix}.$$

So we may take

$$X = \begin{pmatrix} b_{11} & b_{12} & 0 \\ b_{21} & b_{22} & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad Y = t^{-1} \begin{pmatrix} 0 & 0 & b_{13} \\ 0 & 0 & b_{23} \\ -b_{31} & -b_{32} & 0 \end{pmatrix}.$$

(ii) Let first $B \in \mathbb{C}'_{\tilde{A}}$. If $b_{11} = 0$ or $b_{12} \neq b_{31}$, it is enough to find (as above) X and Y such that $B = X + [Y, \tilde{A}]$ and $\tilde{A} + \zeta X \in L_{\tilde{A}}$ for any $\zeta \in \mathbb{C}$. The last condition means that $sp(X) = \{0\}$ and $x_{32} = x_{12}x_{31} = 0$. On the other hand,

$$[Y, \tilde{A}] = \begin{pmatrix} 0 & 0 & y_{12} \\ -y_{31} & -y_{33} & y_{22} - y_{33} \\ 0 & 0 & y_{32} \end{pmatrix}.$$

Let us assume that $b_{31} = 0$ (the computations are similar in the case $b_{12} = 0$). Then we have to choose X of the form

$$X = \begin{pmatrix} b_{11} & b_{12} & b_{13} - y_{12} \\ b_{21} + y_{31} & b_{22} + y_{32} & b_{23} - y_{22} + y_{33} \\ 0 & 0 & -b_{11} - b_{22} - y_{32} \end{pmatrix}$$

such that $\det X = 0$ and $\sigma_2(X) = 0$, that is, $DT = 0$, $D = T^2$, where we write

$$D := \begin{vmatrix} b_{11} & b_{12} \\ b_{21} + y_{31} & b_{22} + y_{32} \end{vmatrix}, \quad T := b_{11} + b_{22} + y_{32}.$$

These conditions are satisfied if and only if

$$y_{32} = -b_{11} - b_{22}, \quad y_{31} = \begin{cases} -b_{21}, & b_{11} = 0 \\ -b_{21} - \frac{b_{11}^2}{b_{12}}, & b_{12} \neq 0 \end{cases}.$$

It remains to show that if $b_{11} \neq 0$ and $b_{12} = b_{31} = 0$, then $k_{\Omega_3}(\tilde{A}; B) > 0$. We may assume that $b_{11} = 1$. Set

$$\tilde{X} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 0 \end{pmatrix}.$$

Choosing X and Y as above, then $B = \tilde{X} + [Y, A_t]$. Let $\alpha > 0$ and $\varphi \in \mathcal{O}(\alpha\mathbb{D}, \Omega_3)$ be such that $\varphi(0) = A_t$ and $\varphi'(0) = B$. Setting $\tilde{\varphi}(\zeta) =$

$e^{-\zeta Y} \varphi(\zeta) e^{\zeta Y}$, then $\tilde{\varphi} \in \mathcal{O}(\alpha \mathbb{D}, \Omega_3)$, $\tilde{\varphi}(0) = \tilde{A}$ and $\tilde{\varphi}'(0) = X$. We get that $k_{\Omega_3}(A_t; B) \geq k_{\Omega_3}(A_t; X)$. The opposite inequality follows in the same way.

Write $\tilde{\varphi}$ in the form

$$\tilde{\varphi}(\zeta) = \tilde{A} + \zeta \tilde{X} + \zeta^2 \hat{X} + o(\zeta^2).$$

Then we compute that

$$\sigma_2(\tilde{\varphi}(\zeta)) = \zeta^2(1 - \hat{x}_{32}) + o(\zeta^2), \quad \det \tilde{\varphi}(\zeta) = -\zeta^3 \hat{x}_{32} + o(\zeta^3).$$

Since $|\sigma_2 \circ \varphi| < 3$, $|\det \varphi| < 1$, we get by the Cauchy inequalities that

$$|\hat{x}_{32} - 1| \leq 3\alpha^{-2}, \quad |\hat{x}_{32}| \leq \alpha^{-3}.$$

So

$$k_{\Omega_3}(A_t; B) = k_{\Omega_3}(\tilde{A}; \tilde{X}) \geq \min_{t \in \mathbb{C}} \max\{\sqrt{|t-1|/3}, \sqrt[3]{|t|}\} > 0.$$

Proof of Corollary 7. Set

$$\tilde{A} = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix}, \quad \tilde{B}_\varepsilon = \begin{pmatrix} 1 & \varepsilon & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 0 \end{pmatrix},$$

$$A = \begin{pmatrix} \tilde{A} & O \\ O & O \end{pmatrix}, \quad B_\varepsilon = \begin{pmatrix} \tilde{B}_\varepsilon & O \\ O & O \end{pmatrix}.$$

It follows as in the proof of Proposition 6 (ii) that

- $k_{\Omega_n}(A; B_0) > 0$;
- for $\varepsilon \neq 0$, there is $\varphi_\varepsilon \in \mathcal{O}(\mathbb{C}, \Omega_n)$ with $\varphi_\varepsilon(0) = A$ and $\varphi'_\varepsilon(0) = B_\varepsilon$.

Then $B_\varepsilon \in C_A$, $\varepsilon \neq 0$, and hence $B_0 \in C_A$.

4. PROOFS OF PROPOSITION 1 AND COROLLARY 2

Proof of Corollary 2. The implication $(a) \Rightarrow (b)$ is trivial and the main implication $(c) \Rightarrow (a)$ is a partial case of Proposition 9 below.

It remains to prove that $(b) \Rightarrow (c)$. Let $\psi(\zeta) = \sum_{k=0}^{\infty} A_k \zeta^k \in \mathcal{O}(s\mathbb{D}, \Omega_n)$ for some $s > 0$. Let $a_{k,\psi} \in \mathbb{C}^{(k+1) \times n^2}$ be the vector with components the entries of A_0, \dots, A_k (taken in some order). Note that

$$\sigma_l(\psi(\zeta)) = \sum_{k=0}^{\infty} p_{l,k}(a_{k,\psi}) \zeta^k, \quad 1 \leq l \leq n,$$

where $p_{j,k}$ are polynomials.

Let now $(r_j) \rightarrow \infty$ and $\varphi_j \in \mathcal{O}(r_j \mathbb{D}, \Omega_n)$, $j \in \mathbb{N}$, uniformly bounded near 0, be such that $\varphi_j(0) = A$ and $\varphi'_j(0) = B$. Then we may assume that $\varphi_j \rightarrow \varphi \in \mathcal{O}(r\mathbb{D}, \Omega_n)$ for some $r > 0$. Hence $p_{l,k}(a_{k,\varphi_j}) \rightarrow p_{l,k}(a_{k,\varphi})$. On the other hand, $|p_{l,k}(a_{k,\varphi_j})| \leq \binom{n}{l} / r_j^k \rightarrow 0$, $k > 0$, by the Cauchy

inequalities. Hence $p_{l,k}(a_{k,\varphi}) = 0$, $k > 0$, that is, $\sigma_l(\varphi(\zeta)) = \sigma_l(\varphi(0)) = \sigma_l(A)$. This means that $\varphi(\zeta) \in L_A$.

Proof of Proposition 1. It is clear that if such a φ exists, then $B \in C_A$.

Conversely, let $B \in C_A$. Then, by [6, p. 86, Proposition 1], there exists a one-dimensional irreducible analytic variety $L_{A,B} \subset L_A$, tangent to B at A . Now, by [6, p. 80, Proposition], there exist $m \in \mathbb{N}$, $r > 0$ and $\psi \in \mathcal{O}(r\mathbb{D}, L_{A,B})$ such that $\psi(\zeta) = A + \zeta^m B + o(\zeta^m)$.

The integer m is the number of sheets in the (local) branched covering provided by the orthogonal projection from L_A to a suitable linear subspace of dimension $n^2 - n$ [6]. This number of sheets corresponds to the cardinality of the solution set, in each generic fiber of the projection, of the equations $\sigma_j(M) = \sigma_j(A)$, $1 \leq j \leq n$. Bezout's theorem shows that this cardinality is less than or equal to the product of the degrees of the polynomials, so here $m \leq n!$.

The above considerations will prove the estimate in Proposition 1, provided that we can replace ψ by an entire map $\tilde{\psi}$ with the same expansion up to order m near 0. So the proof of Proposition 1 reduces to the following.

Proposition 9. *If $A \in \Omega_n$, $m \in \mathbb{N}$ and $\psi \in \mathcal{O}(r\mathbb{D}, L_A)$ for some $r > 0$, then there is $\tilde{\psi} \in \mathcal{O}(\mathbb{C}, L_A)$ with $\tilde{\psi}(\zeta) = \psi(\zeta) + o(\zeta^m)$.*

Proof. We want to reduce that problem by replacing each matrix $\psi(\zeta)$ by a conjugate matrix $\varphi(\zeta)$ (in particular, they will have the same spectrum, so we remain inside L_A and inside Ω_n). If we can manage this so that $\varphi(\zeta)$ is upper triangular, then an entire map with the same spectrum matching φ up to order m can be obtained by taking the Taylor polynomial of degree m of each coefficient of φ .

To proceed with this program, first we need to show that conjugation (with a holomorphic change of basis) does not change the problem.

Let \mathcal{M}_n^{-1} stand for the group of all invertible $n \times n$ matrices.

Lemma 10. *Let $r > 0$, $P \in \mathcal{O}(r\mathbb{D}, \mathcal{M}_n^{-1})$ and $\psi \in \mathcal{O}(r\mathbb{D}, \Omega_n)$.*

Write $\varphi(\zeta) := P(\zeta)^{-1}\psi(\zeta)P(\zeta)$, and assume that there exists $\tilde{\varphi} \in \mathcal{O}(\mathbb{C}, \Omega_n)$ such that near 0, $\tilde{\varphi}(\zeta) = \varphi(\zeta) + o(\zeta^m)$.

Then there exists $\tilde{\psi} \in \mathcal{O}(\mathbb{C}, \Omega_n)$, conjugate to $\tilde{\varphi}$ (in particular they have the same spectrum) such that near 0,

$$\tilde{\psi}(\zeta) = \psi(\zeta) + o(\zeta^m).$$

Note once again that Liouville's theorem implies that the entire maps $\tilde{\varphi}, \tilde{\psi}$ actually map to $L_{\tilde{\varphi}(0)} = L_{\tilde{\psi}(0)}$.

Proof. Note first that, because the exponential is locally Lipschitz, $\exp(A + M) = \exp A + O(M)$.

Denote by $L_m(x)$ the Taylor polynomial of degree m at 0 for the function $x \rightarrow \ln(1 + x)$. Since $\exp(\ln(1 + x)) = 1 + x$ and $\ln(1 + x) = L_m(x) + o(x^m)$, we have $\exp(L_m(x)) = 1 + x + o(x^m)$. So $\exp(L_m(A)) = I + A + o(A^m)$.

Now write

$$P(\zeta) = P(0) \left(I + \sum_{k=1}^m A_k \zeta^k + O(\zeta^{m+1}) \right) =: P(0) (I + M(\zeta)).$$

Define P_1 to be the unique matrix-valued polynomial of degree $\leq m$ in ζ so that

$$L_m \left(\sum_{k=1}^m A_k \zeta^k \right) = P_1(\zeta) + o(\zeta^m).$$

Then, remarking that $M(\zeta) = o(1)$, we have

$$\exp(P_1(\zeta)) = \exp(L_m(M(\zeta)) + o(\zeta^m)) = I + M(\zeta) + o(\zeta^m),$$

so that $P(0) \exp(P_1(\zeta)) = P(\zeta) + o(\zeta^m)$. Then it is easy to see that $\exp(-P_1(\zeta)) P(0)^{-1} = P(\zeta)^{-1} + o(\zeta^m)$, and $\tilde{P}(\zeta) := P(0) \exp(P_1(\zeta))$ defines an entire map. So $\tilde{\psi} := \tilde{P} \tilde{\varphi} \tilde{P}^{-1}$ satisfies the requirements. \square

We now reduce the proof of Proposition 9 to the case of nilpotent matrices (that is, $sp(A) = \{0\}$).

Lemma 11. *Suppose that Proposition 9 holds with the additional hypothesis that $sp(A) = \{0\}$. Then it holds for an arbitrary matrix A .*

Proof. Write $sp(\psi(\zeta)) = sp(A) = \{\mu_j : 1 \leq j \leq k\}$ where the μ_j are distinct eigenvalues with respective algebraic multiplicity m_j . Let $S_j(\zeta) = \ker(\psi(\zeta) - \mu_j I)^{m_j}$ be the associated generalized eigenspace. Choose $\{e_1, \dots, e_n\}$ a basis of \mathbb{C}^n such that $S_j(0) = \text{span}\{e_i : 1 \leq i - \sum_{l=1}^{j-1} m_l \leq m_j\}$. By Lemma 10, without loss of generality we may assume that the matrices are written in this basis, and therefore $\psi(0)$ is a block matrix.

By continuity of the various determinants involved, there exists some $r' > 0$ such that for $|\zeta| < r' \leq r$, we still have, for each j , $\mathbb{C}^n = S_j(\zeta) \oplus \bigoplus_{l:l \neq j} S_l(0)$. Then there is a unique linear projection $\pi_{j,\zeta}$ defined on \mathbb{C}^n such that $\pi_{j,\zeta}(\mathbb{C}^n) = S_j(\zeta)$ and $\text{Ker } \pi_{j,\zeta} = \bigoplus_{l:l \neq j} S_l(0)$. This restricts to a linear isomorphism from $S_j(0)$ to $S_j(\zeta)$. Therefore the vectors $\{\pi_{j,\zeta}(e_i), 1 \leq i - \sum_{l=1}^{j-1} m_l \leq m_j\}$, being obtained as solution of a Cramer system of linear equations with holomorphic coefficients, depend holomorphically on ζ in $D(0, r')$. Thus $\{\pi_{j,\zeta}(e_i), 1 \leq$

$i - \sum_{l=1}^{j-1} m_l \leq m_j, 1 \leq j \leq k\}$ form a basis of \mathbb{C}^n adapted to the direct sum decomposition in the $S_j(\zeta)$. If we write $P(\zeta)$ for the matrix of the coordinates of the vectors of this new basis expressed in the standard basis, it depends holomorphically on ζ in $D(0, r')$ and the new matrix $\hat{\psi}(\zeta) := P(\zeta)^{-1}\psi(\zeta)P(\zeta)$, has the same block structure as $\psi(0)$:

$$\hat{\psi}(\zeta) = \begin{pmatrix} \hat{\psi}_1(\zeta) & 0 & \cdots & 0 \\ 0 & \hat{\psi}_2(\zeta) & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \hat{\psi}_k(\zeta) \end{pmatrix},$$

where $\hat{\psi}_j \in \mathcal{O}(r'\mathbb{D}, \Omega_{m_j})$, $sp(\hat{\psi}_j(\zeta)) = \{\mu_j\}$. The map ω_j defined by

$$\omega_j(\zeta) := \left(\mu_j I_{m_j} - \hat{\psi}_j(\zeta) \right) \left(I_{m_j} - \bar{\mu}_j \hat{\psi}_j(\zeta) \right)^{-1}$$

is in $\mathcal{O}(r'\mathbb{D}, \Omega_{m_j})$, and its values are nilpotent matrices. By our hypothesis there are maps $\tilde{\omega}_j \in \mathcal{O}(\mathbb{C}, \Omega_{m_j})$ such that $\tilde{\omega}_j(\zeta) = \omega_j(\zeta) + o(\zeta^m)$ (and therefore with nilpotent values). Define

$$\tilde{\psi}_j(\zeta) := \left(\mu_j I_{m_j} - \tilde{\omega}_j(\zeta) \right) \left(I_{m_j} - \bar{\mu}_j \tilde{\omega}_j(\zeta) \right)^{-1},$$

and

$$\tilde{\psi}(\zeta) := \begin{pmatrix} \tilde{\psi}_1(\zeta) & 0 & \cdots & 0 \\ 0 & \tilde{\psi}_2(\zeta) & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \tilde{\psi}_k(\zeta) \end{pmatrix}.$$

It is easy to see that $\tilde{\psi} \in \mathcal{O}(\mathbb{C}, \Omega_n)$ and that $\tilde{\psi}(\zeta) = \hat{\psi}(\zeta) + o(\zeta^m)$. \square

Lemma 12. *If $m \in \mathbb{N}$ and $\psi \in \mathcal{O}(r\mathbb{D}, L_0)$ for some $r > 0$, then there is $P \in \mathcal{O}(r'\mathbb{D}, \mathcal{M}_n^{-1})$ such that $\varphi(\zeta) := P(\zeta)^{-1}\psi(\zeta)P(\zeta)$ is a strictly upper triangular matrix for all $\zeta \in r'\mathbb{D}$.*

Proposition 9 follows from Lemma 12.

Indeed, by Lemma 11, we can make the additional hypothesis that ψ has nilpotent values, that is, $\psi \in \mathcal{O}(r\mathbb{D}, L_0)$. By Lemma 12, there is, for every ζ in a neighborhood of 0, a strict upper triangular matrix $\varphi(\zeta) = P(\zeta)^{-1}\psi(\zeta)P(\zeta)$. If we replace each of the holomorphic coefficients $\varphi_{ij}(\zeta)$, $1 \leq i < j \leq n$, by its Taylor polynomial of order m , $\tilde{\varphi}_{ij}(\zeta) := \sum_{k=0}^m \varphi_{ij}^{(k)}(0)\zeta^k/k!$, we obtain $\tilde{\varphi}$ an approximation up to order m of our mapping which is entire, and still strictly upper-triangular, therefore still with spectrum reduced to 0. Since $P(\zeta)$ depends holomorphically on ζ in a neighborhood of 0, we may apply Lemma 10 and obtain the matrix $\tilde{\psi}(\zeta)$ still with spectrum reduced to 0, approximating ψ to order m , and entire in ζ .

Proof of Lemma 12. We are working with a matrix $\psi(\zeta)$ which satisfies $\psi(\zeta)^n = 0$ for all $\zeta \in r\mathbb{D}$. For $1 \leq k \leq n$, let $r_k(\zeta) := \text{rank}(\psi(\zeta)^k)$. For a matrix M , $\text{rank}(M) \leq l < n$ if and only if all the minor determinants of size $l+1$ vanish; in our case those determinants are holomorphic functions of ζ , therefore

$$r_k(\zeta) = \max_{\mathbb{D}} r_k =: \tilde{r}_k$$

for all $\zeta \in r\mathbb{D}$ except on a discrete set. By replacing r by a smaller positive number, if needed, we may assume that $r_k(\zeta) = \tilde{r}_k$ for all $\zeta \in r\mathbb{D}_* := r\mathbb{D} \setminus \{0\}$. Set

$$n_k := n - \tilde{r}_k = \dim \ker \psi(\zeta)^k, \quad \zeta \in r\mathbb{D}_*.$$

It is a classical fact from linear algebra that, for a nilpotent matrix, if $p := \min\{k : n_k = n\}$, then $1 \leq n_1 < \dots < n_p = n_{p+1} = \dots = n$.

For $1 \leq k \leq p$ and $\zeta \in r\mathbb{D}_*$, set $V_k(\zeta) := \ker \psi(\zeta)^k$. Since the Grassmannian $\mathcal{G}(n, n_k)$ is compact, we may find a sequence $(\zeta_i) \rightarrow 0$ and vector subspaces $V_k(0) \in \mathcal{G}(n, n_k)$ such that $\lim_{i \rightarrow \infty} V_k(\zeta_i) = V_k(0) \subset \ker \psi(0)^k$, $1 \leq k \leq p$.

Our problem will be solved if we find $\varepsilon > 0$ and holomorphic mappings $v_j \in \mathcal{O}(\varepsilon\mathbb{D}, \mathbb{C}^n)$ such that $\{v_j(\zeta), 1 \leq j \leq n_k\}$ is a basis of $V_k(\zeta)$, $\zeta \in \varepsilon\mathbb{D}$, $1 \leq k \leq p$. (In particular, $\lim_{\zeta \rightarrow 0} V_k(\zeta) = V_k(0)$).

We shall proceed by induction on k , and on j for each fixed k . The value of ε may be reduced at each step, but we keep the same notation.

By convention we will set $n_0 = 0$, and consider \emptyset as a basis of $\{0\}$. Suppose that we already have determined $\{v_j : 1 \leq j \leq n_k\}$. Choose an $r_{k+1} \times r_{k+1}$ minor matrix of ψ^{k+1} whose determinant, denoted δ_{k+1} , is holomorphic and does not vanish on $\varepsilon\mathbb{D}_*$ and eliminate the unknowns corresponding to the columns of this minor; the other unknowns are then expressed in terms of the former with coefficients which are rational in the coefficients of the matrix ψ^{k+1} , so that we obtain meromorphic vector-valued functions u_i on $\varepsilon\mathbb{D}$ so that $\{u_i(\zeta), 1 \leq i \leq n_{k+1} - n_k\}$ is a basis of $V_{k+1}(\zeta)$ for $\zeta \in \varepsilon\mathbb{D}_*$. Those functions are of the form $u_i := f_i/\delta_{k+1}$, where $f_i \in \mathcal{O}(\varepsilon\mathbb{D}, \mathbb{C}^n)$.

By linear algebra, for each fixed ζ , there exist a set $I(\zeta)$ of $n_{k+1} - n_k$ indices i so that $\{v_j(\zeta), u_i(\zeta) : 1 \leq j \leq n_k, i \in I(\zeta)\}$ form a basis of $V_{k+1}(\zeta)$.

Using the fact that all determinants that we have to compute to determine the rank of a system of vectors are meromorphic in $\varepsilon\mathbb{D}$ and reducing ε if needed, we may choose a fixed set I so that $\{v_j(\zeta), u_i(\zeta) : 1 \leq j \leq n_k, i \in I\}$ form a basis of $V_{k+1}(\zeta)$, for all $\zeta \in \varepsilon\mathbb{D}_*$. Re-index the functions $\{u_i, i \in I\}$ as w_j , $n_k + 1 \leq j \leq n_{k+1}$: we have that

$\{v_j(\zeta), w_l(\zeta) : 1 \leq j \leq n_k < l \leq n_{k+1}\}$ form a basis of $V_{k+1}(\zeta)$ for $\zeta \in \varepsilon\mathbb{D}_*$.

Since δ_{k+1} vanishes at most at 0, we can multiply each vector-valued function w_l by ζ^{α_l} , with $\alpha_l \in \mathbb{Z}$ chosen such that $\zeta^{\alpha_l} w_l(\zeta)$ extends to a map $\tilde{w}_l \in \mathcal{O}(\varepsilon\mathbb{D}, \mathbb{C}^n)$ with $\tilde{w}_l(0) \neq 0$.

We need to modify the \tilde{w}_l to ensure that we still have a system of maximal rank at the origin (this is in the spirit of the Gram-Schmidt orthogonalization process). We proceed by induction on $j \geq n_k + 1$. Suppose we have determined v_j as in the statement of the lemma up to some $j_0 \geq n_k$ such that

$$\begin{aligned} & \text{span}\{v_j(\zeta) : 1 \leq j \leq j_0\} \\ &= \text{span}\{v_j(\zeta), \tilde{w}_l(\zeta) : 1 \leq j \leq n_k < l \leq j_0\}, \quad \zeta \in \varepsilon\mathbb{D}_*, \\ & \text{rank}\{v_j(\zeta) : 1 \leq j \leq j_0\} = j_0, \quad \zeta \in \varepsilon\mathbb{D}. \end{aligned}$$

Let $W(\zeta) := \text{span}\{v_1(\zeta), \dots, v_{j_0}(\zeta), \tilde{w}_{j_0+1}(\zeta)\}$, $\zeta \in \varepsilon\mathbb{D}_*$. Then $\dim W(\zeta) = j_0 + 1$. Again by using the compactness for the Grassmannian, we may choose a sequence $(\zeta_i) \rightarrow 0$ such that $\lim_{i \rightarrow \infty} W(\zeta_i) =: W(0)$ exists. Since all the v_j and \tilde{w}_{j_0+1} are continuous at 0, we easily deduce that $\text{span}\{v_1(0), \dots, v_{j_0}(0), \tilde{w}_{j_0+1}(0)\} \subset W(0)$. Choose a vector w such that $\{v_1(0), \dots, v_{j_0}(0), w\}$ form a basis of $W(0)$.

By reducing ε if needed, we may assume that all the space $W(\zeta)$ and $W(0)$ are each in direct sum with a common supplementary space Y . Then define $v_{j_0+1}(\zeta)$ to be the projection of the fixed vector w to $W(\zeta)$, parallel to Y , for $\zeta \neq 0$; in particular $v_{j_0+1}(0) = w$. Since $v_{j_0+1}(\zeta)$ is obtained by solving a system of linear equations with unique solution, it is meromorphic in a neighborhood of 0, but it is also clearly continuous near 0, hence holomorphic. Since the system $\{v_1(0), \dots, v_{j_0}(0), v_{j_0+1}(0)\}$ is independent, then $\{v_1(\zeta), \dots, v_{j_0}(\zeta), v_{j_0+1}(\zeta)\}$ also is independent for ζ small enough, and since it is contained in $W(\zeta)$ by construction, it forms a basis of that subspace. Reducing ε yet again if needed, the induction may proceed.

REFERENCES

- [1] J. Agler, N. J. Young, *The two-point spectral Nevanlinna-Pick problem*, Integral Equations and Operator Theory 37 (2000), 375–385.
- [2] J. Agler, N. J. Young, *The two-by-two spectral Nevanlinna-Pick problem*, Trans. Amer. Math. Soc. 356 (2004), 573–585.
- [3] J. Agler, N. J. Young, *The hyperbolic geometry of the symmetrized bidisc*, J. Geom. Anal. 14 (2004), 375–403.
- [4] H. Bercovici, C. Foiaş, A. Tannenbaum, *A spectral commutant lifting theorem*, Trans. Amer. Math. Soc. 325 (1991), 741–763.

- [5] R. Brody, *Compact manifolds and hyperbolicity*, Trans. Amer. Math. Soc. 235 (1978), 213–219.
- [6] E. M. Chirka, *Complex analytic sets*, Kluwer Acad. Publ., Dordrecht, Boston, London, 1989.
- [7] C. Costara, *The 2×2 spectral Nevanlinna-Pick problem* J. London Math. Soc. 71 (2005), 684–702.
- [8] C. Costara, *On the spectral Nevanlinna-Pick problem*, Studia Math. 170 (2005), 23–55.
- [9] A. Edigarian, W. Zwonek, *Geometry of the symmetrized polydisc*, Arch. Math. (Basel) 84 (2005), 364–374.
- [10] H.-N. Huang, S. A. M. Marcantognini, N. J. Young, *The spectral Carathéodory-Fejér problem*, Integral Equation and Operator Theory 56 (2006), 229–256.
- [11] M. Jarnicki, P. Pflug, *Invariant distances and metrics in complex analysis—revisited*, Diss. Math. 430 (2005), 1–192.
- [12] N. Nikolov, *Continuity and boundary behavior of the Caratheodory metric*, Math. Notes, 67 (2000), 183–191.
- [13] N. Nikolov, P. J. Thomas, W. Zwonek, *Discontinuity of the Lempert function and the Kobayashi-Royden metric of the spectral ball*, Preprint, 2007 (arXiv:math.CV/0704.2470).
- [14] J. Yu, *Singular Kobayashi metrics and finite type conditions*, Proc. Amer. Math. Soc. 123 (1995), 121–130.

INSTITUTE OF MATHEMATICS AND INFORMATICS, BULGARIAN ACADEMY OF SCIENCES, ACAD. G. BONCHEV 8, 1113 SOFIA, BULGARIA
E-mail address: `nik@math.bas.bg`

LABORATOIRE EMILE PICARD, UMR CNRS 5580, UNIVERSITÉ PAUL SABATIER, 118 ROUTE DE NARBONNE, F-31062 TOULOUSE CEDEX, FRANCE
E-mail address: `pthomas@cict.fr`